

Dynamical switching of networks facilitates endemicity in the susceptible-infected-susceptible epidemic model

Leo Speidel¹, Konstantin Klemm^{2,3}, Víctor M. Eguíluz³, and Naoki Masuda^{*4}

¹ Doctoral Training Centre in Systems Biology, University of Oxford, Rex Richards Building, South Parks Road, Oxford OX1 3QU, United Kingdom

² School of Science and Technology, Nazarbayev University, Qabanbay Batyr Ave 53, Astana 010000, Kazakhstan

³ Instituto de Física Interdisciplinar y Sistemas Complejos IFISC (CSIC-UIB), E07122 Palma de Mallorca, Spain

⁴ Department of Engineering Mathematics, University of Bristol, Merchant Venturers Building, Woodland Road, Clifton, Bristol BS8 1UB, United Kingdom

E-mail: *naoki.masuda@bristol.ac.uk

Keyword: temporal networks, epidemic dynamics, SIS model, epidemic threshold

Abstract

Various data recorded from humans, animals, and computers suggest that networks underlying epidemic processes are time-varying. Epidemic processes on temporal networks are complicated by complexity of time-varying networks. In particular, how the two timescales, one from that of an epidemic process and the other from the dynamics of network structure, interplay and potentially change the traditional understanding of epidemic processes on networks, which is mostly based on static networks, has to be revealed [1]. In this presentation, we report a theory to understand the susceptible-infected-susceptible epidemic model on arbitrary temporal networks [2]. We show that, under our modelling framework, temporality of networks always lessens the epidemic threshold such that infections persist more easily in temporal networks than in their static counterparts. We also propose a quantity to account for why the epidemic threshold is substantially lowered in some temporal networks than others.

References

- [1] N. Masuda and P. Holme, “Predicting and controlling infectious disease epidemics using temporal networks”, F1000Prime Reports, vol. 5, article no. 6, (2013).
- [2] L. Speidel, K. Klemm, V. M. Eguíluz, and N. Masuda, “Temporal interactions facilitate endemicity in the susceptible-infected-susceptible epidemic model”, Preprint: arXiv:1602.00859 (2016).